

Non-Parallel Electric and Magnetic Fields in a Gravitational Background, Stationary Gravitational Waves and Gravitons

*Carlos Pinheiro - **J.A. Helayël Neto - ***Gilmar S. Dias
(*Departamento de Física, CCE, UFES. **Centro Brasileiro
de Pesquisas Físicas, CBPF. Univesidade Católica de Petró-
polis, UCP. ***Departamento de Física, CCE,
UFES. Escola Técnica Federal ETFES)

Abstract

The existence of an electromagnetic field with parallel electric and magnetic components is readdressed in the presence of a gravitational field. A non-parallel solution is shown to exist. Next, we analyse the possibility of finding stationary gravitational waves in de nature. Finaly, We construct a D=4 effecttive quantum gravity model. Tree-level unitarity is verified.

1 Electric and Magnetic Field in a Gravitational Background

Based on a series of papers by Brownstein [1] and Salingaros [2], we readdress here the possibility of the existence of an electromagnetic field whose electric and magnetic components are parallel in the presence of a gravitational field. The coupling between the electromagnetic sector and the gravitational background is accomplished by means of the action.

$$\mathcal{S} = \int \sqrt{-\tilde{g}} \left(-\frac{1}{4} F_{\mu\nu} F^{\mu\nu} \right) d^4x, \quad (1.1)$$

where

$$\tilde{g} = \det(g_{\mu\nu}),$$

and

$$F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu.$$

From the above action, the following field-equations follow:

$$\mathcal{D}_\mu F^{\mu\nu} = \partial_\mu F^{\mu\nu} + \Gamma_{\beta\lambda}^\beta F^{\lambda\nu} + \Gamma_{\mu\lambda}^\nu F^{\mu\lambda} = J^\nu \quad (1.2)$$

$$\mathcal{D}_\mu F_{\nu\beta} + \mathcal{D}_\nu F_{\beta\mu} + \mathcal{D}_\beta F_{\mu\nu} = 0. \quad (1.3)$$

Choosing the background to be described by the F.R.W metric,

$$dS^2 = dt^2 - a^2(t) \left[\frac{dr^2}{1 - Ar^2} + r^2 d\theta^2 + r^2 \sin^2 \theta d\phi^2 \right], \quad (1.4)$$

the Maxwell equations in the absence of electromagnetic sources read as below:

$$\begin{aligned} \vec{\nabla} \cdot \vec{E} &= g \vec{\nabla} f \cdot \vec{E} \\ \vec{\nabla} \cdot \vec{B} &= 0 \\ \vec{\nabla} \times \vec{E} &= -\frac{\partial \vec{B}}{\partial t} \\ \vec{\nabla} \times \vec{B} &= \frac{\partial \vec{E}}{\partial t} - g \frac{\partial f}{\partial t} \vec{E} + g \vec{\nabla} f \times \vec{B} - \Gamma_{\mu\beta}^i F^{\mu\beta}, \end{aligned} \quad (1.5)$$

where

$$g = \frac{\sqrt{1 - Ar^2}}{a^3 r^2 \sin \theta}, \quad f = \frac{a^3 r^2 \sin \theta}{\sqrt{1 - Ar^2}}, \quad (1.6)$$

and $A = +1, 0, -1$.

The Wave-equations for \vec{E} and \vec{B} are found to be given by:

$$\nabla^2 \vec{E} - \frac{\partial^2 \vec{E}}{\partial t^2} = \vec{\nabla}(g \vec{\nabla} f \cdot \vec{E}) - \frac{\partial g}{\partial t} \frac{\partial f}{\partial t} \vec{E} - g \frac{\partial^2 f}{\partial t^2} \vec{E} - g \frac{\partial f}{\partial t} \frac{\partial \vec{E}}{\partial t} + \quad (1.7)$$

$$+ \frac{\partial g}{\partial t} \vec{\nabla} f \times \vec{B} + g \frac{\partial}{\partial t} \vec{\nabla} f \times \vec{B} + g \vec{\nabla} f \times \frac{\partial \vec{B}}{\partial t} - \frac{\partial}{\partial t} (\Gamma_{\mu\beta}^i F^{\mu\beta});$$

$$\nabla^2 \vec{B} - \frac{\partial^2 \vec{B}}{\partial t^2} = \vec{\nabla} \times (g \frac{\partial f}{\partial t} \vec{E} - g \vec{\nabla} f \times \vec{B} + \Gamma_{\mu\beta}^i F^{\mu\beta}) \quad (1.8)$$

Now, by virtue of the presence of the gravitational background, we have explicitly built up a solution for \vec{E} and \vec{B} that evade the claim by Brownstein [1] and Salingaros [2]. These authors state that it is always possible to find out parallel solutions for \vec{E} and \vec{B} in Plasma Physics or in an Astrophysic Plasma. However, contrary to their result, we have found non-parallel solutions due to the non-flat background of gravity:

$$\begin{aligned} \vec{E} = & \hat{i}(\sin \theta G_{(r,t,\theta)} - \cos \theta F_{(r,t)}) ka \cos(kz) \cos(\omega t) + \\ & + \hat{j}(\cos \theta F_{(r,t)} - \sin \theta G_{(r,t,\theta)}) ka \sin(kz) \cos(\omega t) + \\ & + \hat{k}[(\sin \theta \cos \varphi F_{(r,t)} + \cos \theta \cos \varphi G_{(r,t,\theta)}) ka \cos(kz) \cos(\omega t) + \\ & - (\sin \theta \sin \varphi F_{(r,t)} + \cos \theta \sin \varphi G_{(r,t,\theta)}) ka \sin(kz) \cos(\omega t)] \end{aligned} \quad (1.9)$$

and

$$\vec{B} = ka[\hat{i} \sin(kz) + \hat{j} \cos(kz)] \cos(\omega t) \quad (1.10)$$

where the functions $G_{(r,t,\theta)} = \frac{a \cot g \theta}{3\dot{a}r}$ and

$$F_{(r,t)} = \frac{2a}{3\dot{a}r} + \frac{Aar}{3\dot{a}(1 - Ar^2)} \quad (1.11)$$

are the metric contribution.

2 Stationary Gravitational Waves and Gravitons

Now, we analyse the possibility of finding stationary gravitational waves. From a phenomenological viewpoint, a distribution of black holes could play the role of knots for the non-propagating gravitational waves. We postulate the equation that may lead to this sort of waves:

$$R_{\mu\nu} = \kappa\Lambda h_{\mu\nu}, \quad (2.12)$$

$$g_{\mu\nu}(x) = \eta_{\mu\nu} + \kappa h_{\mu\nu}, \quad (2.13)$$

where Λ is the cosmological constant. These equations yield:

$$\partial_\beta \partial_\nu h_\mu^\beta + \partial_\beta \partial_\mu h_\nu^\beta - \square h_{\mu\nu} - \partial_\mu \partial_\nu h_\beta^\beta = \Lambda h_{\mu\nu}. \quad (2.14)$$

Now, solution of the form

$$h_{\mu\nu} = C_{\mu\nu}(z)f(t), \quad (2.15)$$

$$h_{\mu\nu} = \begin{pmatrix} A_{00} & 0 & 0 & 0 \\ 0 & A_{11} & A_{12} & 0 \\ 0 & A_{12} & -A_{11} & 0 \\ 0 & 0 & 0 & A_{00} \end{pmatrix} e^{i\tilde{k}z} \cos \omega t, \quad (2.16)$$

can be found, where A_{00} , A_{11} and A_{12} are free parameters, whereas $\tilde{k} = \sqrt{\Lambda - \omega^2}$ is the wave number. Having in mind that Λ is a small number, the frequency ω must be extremely small. This forces on us to search for a mechanism to detect such low-frequency stationary waves.

The equations of motion stemming from 2.12 may be derived from the density Lagrangian

$$\begin{aligned} \mathcal{L}_H = & \frac{1}{2} H^{\mu\nu} \square H_{\mu\nu} - \frac{1}{4} H \square H - \frac{1}{2} H^{\mu\nu} \partial_\mu \partial_\alpha H_\nu^\alpha - \frac{1}{2} H^{\mu\nu} \partial_\nu \partial_\alpha H_\mu^\alpha + \\ & - \frac{1}{2} \Lambda H^{\mu\nu} H_{\mu\nu} + \frac{1}{4} \Lambda H^2, \end{aligned} \quad (2.17)$$

where

$$H_\nu^\alpha = h_\nu^\alpha - \frac{1}{2}\delta_\nu^\alpha h$$

and the bilinear form correspondent operator of Lagrangian 2.17 is given by

$$\begin{aligned} \Theta_{\mu\nu,\kappa\Lambda} = & (\square - \Lambda)P^{(2)} - \Lambda P_m^{(1)} + \frac{5}{2}(\square - \Lambda)P_s^{(0)} - \frac{(\Lambda + 3\square)}{2}P_w^{(0)} + \\ & + \frac{\sqrt{3}}{2}(\Lambda - \square)P_{sw}^{(0)} + \frac{\sqrt{3}}{2}(\Lambda - \square)P_{ws}^{(0)}, \end{aligned} \quad (2.18)$$

and $P^{(i)}$, $i=0,1,2$, are spin-projetor operators in the space of rank-2 symmetric tensors. The graviton propagator read off from this Lagrangian is given by:

$$\langle T(h_{\mu\nu}(x); h_{\kappa\lambda}(y)) \rangle = i\Theta_{\mu\nu,\kappa\lambda}^{-1}\delta^4(x - y) \quad (2.19)$$

where

$$\Theta^{-1} = [XP^{(2)} + YP_m^{(1)} + ZP_s^{(0)} + WP_w^{(0)} + RP_{sw}^{(0)} + SP_{ws}^{(0)}]_{\mu\nu,\kappa\lambda} \quad (2.20)$$

with

$$X = -\frac{1}{\Lambda - \square}; \quad Y = -\frac{1}{\Lambda}; \quad Z = -\frac{\Lambda + 3\square}{\Lambda^2 + 8\Lambda\square - 9\square^2}; \quad (2.21)$$

$$W = -\frac{5}{\Lambda - 9\square}; \quad R = -\frac{\sqrt{3}}{\Lambda + 9\square}; \quad S = -\frac{\sqrt{3}}{\Lambda + 9\square};$$

From this propagator, we can set a current-current amplitude and discuss tree-level unitarity [3]. Three massive excitations were found: They are a spin-2 quantum with mass equal to $k^2 = \Lambda$ and two massive spin-0 quanta with masses equal to $k^2 = \Lambda$ and $k^2 = -\frac{1}{9}\Lambda$. The spin-2 is a physical one: the imaginary part of the residue of the amplitude at the pole $k^2 = \Lambda$ is positive, so that it does not lead to a ghost. It remains to be shown that the tachyonic pole, $k^2 = -\frac{1}{9}\Lambda$, is non-dynamical or decouples through some constraint on the sources.

We conclude, then, that in a gravitational background it is always possible to find non-parallel electric and magnetic fields. It is the gravitational field that breaks the parallel configuration of \vec{E} and \vec{B} argued by [1, 2]. Also, a stationary gravitational wave equation was postulated and a particular solution was found. We argue that such a kind of solution is possible to be found in Black Hole distributions. Finally we set up an effective quantum gravity model where the necessary condition for the tree-level unitarity for the spin-2 sector is respected. The model is infrared finite though non-renormalizable in the ultraviolet limit.

Acknowledgements

The authors acknowledge Dr. Berth Schröer for helpful suggestions and technical discussions. Thanks are also due to Gentil O. Pires and Manoelito M. de Souza for a critical reading of the manuscript. This work was partially supported by the Conselho de Desenvolvimento Científico e Tecnológico - CNPQ - Brazil.

References

- [1] K.R.Brownstein, Phys. Rev A 35 (1987) - 4854;
- [2] N.Salingaros, Phys. Rev A 19 (1986) L 101; Am. J. Phys. 53, 361 (1985);
- [3] C. Pinheiro, G.O.Pires, Phys. Lett. B 301 (1993) 339; “Some Quantum Aspects of D=3 Space-Time Massive Gravity” preprint IF-UFES; C.Pinheiro, G.O.Pires, N.Tomimura, “Some Quantum Aspects of Three-Dimensional Einstein-Chern-Simons-Proca Massive Gravity”, Il Nuovo Cimento, to appear; C.Pinheiro, G.O.Pires, C.Sasaki, “On a Three-Dimensional Gravity Model With Higher Derivatives”, General Relativity and Gravitation, to appear.